

Code : 103102

B.Tech 1st Semester Exam., 2019  
(New Course)

MATHEMATICS—I

( Calculus and Differential Equations )

Time : 3 hours

Full Marks : 70

Instructions :

- (i) The marks are indicated in the right-hand margin.
- (ii) There are **NINE** questions in this paper.
- (iii) Attempt **FIVE** questions in all.
- (iv) Question No. 1 is compulsory.

1. Choose the correct answer of the following  
(any seven) : 2×7=14

(a) The minimum value of the function

$$f(x) = \sin x(1 + \cos x), 0 < x < 2\pi$$

is

- (i)  $\pi$
- (ii)  $\frac{2\pi}{3}$
- (iii)  $\frac{5\pi}{3}$
- (iv)  $2\pi$

20AK/276

( Turn Over )

(b) If

$$f(a+b-x) = f(x), \text{ then } \int_a^b x f(x) dx$$

is equal to

- (i)  $\left(\frac{a+b}{2}\right) \int_a^b f(b-x) dx$
- (ii)  $\left(\frac{a+b}{2}\right) \int_a^b f(x) dx$
- (iii)  $\left(\frac{b-a}{2}\right) \int_a^b f(x) dx$
- (iv)  $\left(\frac{a-b}{2}\right) \int_a^b f(x) dx$

(c) The slope of the tangent to the curve

$$y = \int_0^{x^2} \left( \frac{dt}{1+t^3} \right)$$

at the point where  $x=1$ , is

- (i) 2
- (ii) 1
- (iii)  $\frac{1}{2}$
- (iv)  $\frac{1}{4}$

(d) The value of

$$\lim_{x \rightarrow 0} \frac{xe^{x^2}}{\int_0^x e^{t^2} dt}$$

is

- (i) 0
- (ii) 1
- (iii) 2
- (iv) -1

(e) The series whose  $n$ th term is

$$\{(n^3 + 1)^{\frac{1}{3}} - n\}$$

is

- (i) convergent
- ~~(ii) divergent~~
- (iii) oscillatory
- (iv) None of the above

(f) The directional derivative of

$$\varphi(x, y, z) = x^2yz + 4xz^2$$

at the point  $(1, -2, -1)$  in the direction  $2i - j - 2k$  is

- (i) 1
- (ii) 3
- (iii)  $\frac{11}{3}$
- ~~(iv)  $\frac{37}{3}$~~

(g) The general solution of PDE

$$uu_x + yu_y = x$$

is

(i)  $u^2 = g\left(\frac{y}{x+u}\right) + x^2$

(ii)  $f(u^2 + x^2) = 0$

~~(iii)  $f(x+y) = 0$~~

(iv) None of the above

(h) If  $J_n$  is the Bessel's function of first kind, then the value of  $J_{\frac{3}{2}}$  is

(i)  $\sqrt{\frac{2}{\pi x}} \left( \frac{\cos x}{x} - \sin x \right)$

(ii)  $\sqrt{\frac{2}{\pi x}} \left( \frac{\sin x}{x} - \cos x \right)$

(iii)  $\sqrt{\frac{2}{\pi x}} \sin x$

~~(iv)  $\sqrt{\frac{2}{\pi x}} \cos x$~~

http://www.akubihar.com

http://www.akubihar.com

http://www.akubihar.com

http://www.akubihar.com

(i) The general solution of

$$\frac{d^2y}{dx^2} + 9y = \sin^3 x$$

is

~~(i)~~  $y = A \cos(3x + B) + \frac{1}{24} \sin x - \sin 3x$

(ii)  $y = Ae^{3x} + Be^{-3x} + \frac{1}{32} \sin x + \frac{1}{2} \cos 3x$

(iii)  $y = A + Be^{3x} + 2 \sin x - \frac{5}{13} \sin 3x$

(iv)  $y = A \sin(3x + B) + \frac{3}{32} \sin x + \frac{x}{24} \cos 3x$

(j) If  $P_n$  is the Legendre polynomial of first kind, then the value of

$$\int_{-1}^1 P_{n+1}^2 dx$$

is

~~(i)~~  $\frac{2}{(2n+1)}$

(ii)  $\frac{2}{(2n+2)}$

(iii)  $\frac{2}{(2n+3)}$

(iv)  $\frac{2}{(2n+4)}$

2. (a) Find the evolutes of the curve

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

where  $a > b$ .

(b) Prove that  $\Gamma(1/2) = \sqrt{\pi}$ .

7+7

3. (a) Find the extreme values of

$$f(x, y, z) = 2x + 3y + z$$

such that  $x^2 + y^2 = 5$  and  $x + z = 1$ .

(b) Find  $\theta$ , if

$$f(x+h) = f(x) + hf'(x) + \frac{h^2}{2!} f''(x+\theta h)$$

$$0 < \theta < 1 \text{ and } f(x) = ax^3 + bx^2 + cx + d. \quad 7+7$$

4. (a) Discuss the continuity of the function

$$f(x, y) = \begin{cases} \frac{x-y}{x+y}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

(b) Expand in the sine series of the function

$$f(x) = \begin{cases} x, & 0 \leq x \leq 4 \\ 8-x, & 4 \leq x \leq 8 \end{cases}$$

7+7

5. (a) Find the volume of the solid generated by revolving the ellipse  $4x^2 + 9y^2 = 36$ .

(b) Evaluate the integral by changing to polar coordinates

$$\int_0^1 \int_0^{\sqrt{2x-x^2}} (x^2 + y^2) dy dx \quad 7+7$$

6. (a) Find the mass of a plate in the first quadrant of an ellipse  $2x^2 + 3y^2 = 1$ , whose density per unit area is given by  $\rho = kxy$ .

(b) Find the directional derivative of  $\phi(x, y, z) = x^2yz + 4xz^2$  at  $(1, -2, -1)$  in the direction  $2i - j - 2k$ . 7+7

7. (a) Evaluate  $\oint_C (xy) dx + (x^2 + y^2) dy$ , around the boundary of the region defined by  $y^2 = 8x$  and  $x = 2$ , using Green's theorem.

(b) Find  $(\nabla \times A) \times B$ , where

$$\vec{A} = x^2zi + yz^3j - 3xyk \text{ and}$$

$$\vec{B} = 3xi + 4zj - xyk$$

7+7

( Turn Over )

8. (a) Solve the differential equation

$$\frac{d^2y}{dx^2} + 9y = \sec 3t$$

by variation of parameters.

(b) Solve the differential equation

$$1 + y^2 + (x - e^{-\tan^{-1}y}) \frac{dy}{dx} = 0. \quad 7+7$$

9. (a) Prove that

$$\int_{-1}^1 x^2 P_{n+1} P_{n-1} dx = \frac{2n(n+1)}{(2n-1)(2n+1)(2n+3)}$$

(b) Find the complete integral of the partial differential equation

$$2xz + q^2 = x(px + qy) \quad 5+9$$